



Fact Sheet

OTITIS MEDIA

What is Otitis Media? Otitis media is another name for an ear infection. It is one of the most common problems for which a physician sees a child. It is an inflammation of the middle ear space and is usually accompanied by fluid build-up. This fluid may or may not be infected. This build up of fluid in the middle ear space restricts the movement of the ear drum. If the ear drum does not move freely, a hearing loss occurs. This can be compared to plugging your ear with your finger. The child with otitis media does not always show signs such as fever, irritability, tender ears, reddened ears, etc.

There are three general categories of otitis media:

1. otitis media without effusion (fluid)
2. otitis media with effusion
3. otitis media with perforation (hole in ear drum)

What causes it? The Eustachian tube is the tube connecting the middle ear and the upper part of the nasal passages. Eustachian tube dysfunction is a significant factor in the development of otitis media. The purpose of the Eustachian tube is to equalize the pressure on both sides of the eardrum and provide ventilation to the middle ear space. When it does not work properly, the air is trapped in the middle ear space. The air is absorbed and fluid results.

Age plays a factor in the development of otitis media. At birth, the Eustachian tube is in a horizontal position and it is wider and shorter. During the first few years of life it begins to extend downward allowing for easier drainage. Therefore, otitis media is most common during the first 2 years of life. Inflammations of the nasal end of Eustachian tube may produce swelling, thus impairing its function. Such inflammation may result from viral or bacterial infection (a cold) or chemical irritation (tobacco smoke, chlorinated pool water). Allergies have also been known to cause otitis media. Some foods that commonly cause allergic reactions in young children include milk, wheat, eggs, corn, yeast, peanuts, soybean, sugar and citrus.

Skeletal changes play an important part in the development of otitis media. Children with Down syndrome, cranial base problems (cleft palate, atresia, adenoid problems, etc.) and craniofacial syndrome (Treacher-Collins, Crouzon's or CHARGE Association) have a high incidence of otitis media. Children with upper respiratory problems are also more prone to develop otitis media.

Children with upper respiratory problems are also more prone to develop otitis media. Family history, climate, and dietary reasons have also been linked to recurring otitis media.

What are the effects? Recurring otitis media has been shown to have a direct impact on a child's speech, language, and academic development. If a hearing loss is present due to otitis media, a child will experience difficulties in attending and following both formal communication (classroom instruction), and incidental communication (conversation that is on-going throughout the day).

Amplification (e.g., hearing aids) may not be prescribed for the child with otitis media because in the majority of cases, the child's hearing returns to normal after the otitis media is cleared. However, more severe cases, especially when prolonged or left untreated, may result in a perforated ear drum, scar tissue on the ear drum and even a permanent hearing loss. Other, more serious, complications that could occur include facial nerve paralysis, meningitis, encephalitis or brain abscess.

Children that have a known sensorineural hearing loss may experience a decrease in their usable hearing and, in some cases; these children may not be able to wear their hearing aids during an episode of otitis media. The ear must be allowed to breathe and drain. If an ear mold or hearing aid is in the ear, increased condensation may occur or the fluid may drain into the hearing aid causing internal damage to the amplifier. Some children may experience such pain that they cannot tolerate the ear mold or hearing aid in their ear.

What is the treatment? Treatment for otitis media varies according to the severity, the occurrence, and the age of the child. If the otitis media is found to be infectious, antibiotics are usually recommended. Clearance of the liquid is imperative. If the child has a history of recurring otitis media the doctor may choose to perform a myringotomy and place a tympanostomy tube in the ear drum. This is a common procedure in young children that entails the doctor making an incision in the ear drum to drain the fluid. A small tube is then placed in the incision. This tube allows for the air pressure to be equalized on both sides. It also assists in keeping the middle ear space well ventilated.

If you suspect your child may be prone to otitis media consult your physician, pediatrician or ear, nose, and throat specialist (ENT).

GLOSSARY

effusion – The escape of fluid into the middle ear cavity.

perforation – A hole or tear

cranial base anomalies – Pertaining to or involving both the braincase and face.

cleft palate – Nonclosure of all or part of the roof of the mouth.

atresia – Absence of the ear canal.

myringotomy – A small cut in the ear drum.

facial paralysis – Paralyzed face muscles.

meningitis – Inflammation of the membranes covering the brain and spinal cord.

encephalitis – Inflammation of the brain.

For more information contact:

Tanni Anthony
Colorado Department of Education
Special Education Services Unit
201 E. Colfax Avenue, Suite 300
Denver, CO 80203
Phone: 303-866-6681
Anthony_T@cde.state.co.us

or

Gina Quintana
Colorado Department of Education
Special Education Services Unit
201 E. Colfax Avenue, Suite 300
Denver, CO 80203
303-866-6605
Quintana_G@cde.state.co.us

Fax: 303-866-6811

TTY: 303-860-7060

Fact Sheets from the Colorado Services to Children with Deafblindness are to be used by both families and professionals serving individuals with vision and hearing loss. The information applies to children, birth through 21 years of age. The purpose of the Fact Sheet is to give general information on a specific topic. More specific information for an individual student can be provided through personalized technical assistance available from the Colorado Deafblindness Project. For more information call (303) 866-6681 or (303) 866-6605.

Information for the Fact Sheet kindly supplied by California Deaf-Blind Services – January 2002