



Fact Sheet

Assessments for Hearing & Functioning of the Ear

By Jill Grattan & MaryAnn Demchak
Nevada Dual Sensory Impairment Project

Hearing assessments attempt to determine the degree of hearing loss (mild, moderate, profound), type of hearing loss (see below), and configuration of hearing loss (e.g., bilateral or loss in both ears, unilateral or loss in one ear, etc.). Other tests determine how parts of the ear are functioning and are not actual tests of hearing.


Definitions


Physiological Testing: Objective tests or measures that rely on recorded responses from the body; the individual being tested does not have to respond as is required for behavioral testing


Behavioral Testing: Tests that require a response from the individual (e.g., raising one's hand in response to a tone, repeating a word presented through earphones)






Image of Tympanometry

Category	Test name	What it tests	How the test is conducted	Age testing can be conducted
Physiological Testing <i>Image of Tympanometry</i> 	Tympanometry	<ul style="list-style-type: none"> • Purpose is to evaluate the function of the middle ear and tympanic membrane ⁴ • Does not tell if child is hearing or not ³ • Results reveal how well the middle ear is functioning ⁴ • When combined with Acoustic Reflex Test (below), the results reveal how well the middle ear is functioning ⁴ 	<ul style="list-style-type: none"> • Non-invasive test • Takes approximately two minutes to perform ⁴ • A probe is inserted into each of the patient's ears. • Results are not dependent on a response from the patient; however, the patient should not move ⁴ 	Can be performed on infants from 7 months of age through adults ⁶
	Acoustic Reflex Test	<ul style="list-style-type: none"> • Purpose is to measure middle ear muscle in response to sounds ⁶ • The muscle in the middle ear contracts as a reflex in response to sounds • When combined with tympanometry (above), the results reveal how well the middle ear is functioning ⁴. 	<ul style="list-style-type: none"> • A probe is inserted into each of the patient's ears. • The results are not dependent on a response from the patient; however, the patient should not move ⁴ 	Can be performed on infants from 7 months of age through adults ⁶

<p style="writing-mode: vertical-rl; transform: rotate(180deg);">Image of OAE testing</p> 	<p>Otoacoustic Emissions Testing (OAEs)</p>	<ul style="list-style-type: none"> • Test to determine how well the inner ear functions, specifically, the cochlea ⁷. • Otoacoustic emissions are sounds from vibrations produced by the outer hair cells of the cochlea (in the inner ear); these hair cells vibrate when the cochlea is stimulated by sound • Test can, "partially estimate hearing sensitivity within a limited range ..." ⁷ • Individuals with a hearing loss greater than 25-30 dB will not produce these otoacoustic emissions 	<ul style="list-style-type: none"> • A small plug is inserted into the patient's ear and a microphone in the plug records responses of the ear ⁷. • This test can be conducted while the patient is asleep, takes a few minutes ⁷ 	<p>Can be performed on newborns through adults ⁷</p>
---	--	--	---	---

Category	Test name	What it tests	How the test is conducted	Age testing can be conducted
Physiological Tests	Auditory Brainstem Response (ABR)	<ul style="list-style-type: none"> • Gives information about inner ear and brain pathways for hearing • The purpose of this test is to measure the function of the brainstem in response to sounds ⁵ 	<ul style="list-style-type: none"> • Earphones are placed into the patient's ears and electrodes are placed onto the patient's head ³ • Clicking noises are sent through the earphones and the electrodes measure the brain's activity ³ • No response is needed from the person • Takes a few minutes and can be conducted while the patient is sleeping ³ 	Can be performed on newborns through adults ⁵
<p>Images of ABR testing</p> 				

Category	Test name	What it tests	How the test is conducted	Age testing can be conducted
Behavioral Tests	Behavioral Audiometry	<ul style="list-style-type: none"> • This test is used for screening purposes ³ 	<ul style="list-style-type: none"> • Observing infant's behavior in response to certain sounds is observed ³ 	Can be performed on infants through adults ³
	Pure-Tone Audiometry Or Pure-Tone Air Conduction Testing	<ul style="list-style-type: none"> • Test measures hearing sensitivity, specifically, the softest sound an individual can hear at specific frequencies 50% of the time ⁹ • "This test assesses sensitivity when the signal is transmitted through the outer, middle, and inner ear, through the brain to the cortex" ¹⁰ 	<ul style="list-style-type: none"> • The patient may wear earphones or headphones or test may be conducted using speakers ¹⁰ • When earphones are worn, results for each ear are obtained • If test is done using speakers, it is not possible to obtain ear-specific results • The person is asked to make a response (e.g., raise hand) when he/she hears a sound ³ 	Can be performed to assess children older than 4 years old ³
	Pure-Tone Bone Conduction Testing	<ul style="list-style-type: none"> • The purpose of this test is to test hearing in the inner ear without utilizing the outer or middle ear ^{10, 1} • Used if there is blockage in outer or middle ear 	<ul style="list-style-type: none"> • A small vibrator is placed behind the ear or on the forehead of the patient ¹. • The signal gently vibrates the bones of the skull, and directly stimulates the inner ear ¹ 	Can be performed on infants through adults ⁹
<p>Image of Bone Conduction Testing</p>  <p>Image of Behavioral Audiometry Testing</p> 				

<i>Category</i>	<i>Test name</i>	<i>What it tests</i>	<i>How the test is conducted</i>	<i>Age testing can be conducted</i>	
Behavioral Tests	Visual Reinforcement Audiometry (VRA)	<ul style="list-style-type: none"> Variation on the Pure-Tone Audiometry test (above)¹ "This test assesses sensitivity when the signal is transmitted through the outer, middle, and inner ear, through the brain to the cortex"¹⁰. 	<ul style="list-style-type: none"> Patients are tested wearing earphones or headphones or in a sound booth with speakers¹ The patient is taught to look at a visual object (e.g., flashing light, moving toy) when a sound is heard¹ 	Can be performed to assess children from 6 months to adult ³ . Typical ages: 6 months to 2 years old ¹	
		Conditioned Play Audiometry (image above)	<ul style="list-style-type: none"> Variation of the Pure-Tone Audiometry test (above)¹ "This test assesses sensitivity when the signal is transmitted through the outer, middle, and inner ear, through the brain to the cortex"¹⁰ 	<ul style="list-style-type: none"> Patients are tested wearing earphones or headphones or in a sound booth with speakers¹ The patient is taught to perform a task (e.g., drop a block into a box, stack a ring) each time a sound is heard¹ 	Can be performed to assess children from 2 years to adult ¹ . Typical ages: 2 to 5 years old ¹
	Speech Awareness Threshold (SAT) Or Speech Detection Threshold (SDT)	<ul style="list-style-type: none"> Indicates the lowest level at which speech can be detected at least 50% of the time⁸ 	<ul style="list-style-type: none"> Patient listens to a voice say words via earphones or loud speakers⁸ and indicates when speech is present⁸ 	Can be performed to assess children who are too young to repeat ⁸	
	Speech Reception Threshold or Speech Recognition Threshold (SRT)	<ul style="list-style-type: none"> Indicates the lowest level at which speech can be identified at least 50% of the time⁸ 	<ul style="list-style-type: none"> Patient listens to a voice say words via earphones or loud speakers² and repeats word he/she heard or indicates word recognition² 	Can be performed to assess older children and adults ²	

References

- American Speech-Language-Hearing Association (n.d). *Pure-Tone Testing*. Retrieved December 21, 2010, from <http://www.asha.org/public/hearing/Pure-Tone-Testing/>
- American Speech-Language-Hearing Association (n.d). *Speech Testing*. Retrieved December 21, 2010, from <http://www.asha.org/public/hearing/Speech-Testing/>
- Children's Hospital of Wisconsin, 2010. (n.d.). *Types of Hearing Tests*. Retrieved December 21, 2010, from <http://www.chw.org/display/PPF/DocID/21456/router.asp>
- Mikolai, T. K., Duffey, J., Adlin, D. from Maico Diagnostics. (2006). *A guide to tympanometry for hearing screening*. Retrieved December 23, 2010, from http://www.maico-diagnostics.com/eprise/main/_downloads/com_en/Documentation/Guide.Tymp.pdf
- Bhattacharyya, N. Emedicine from *WebMD*. (2009, January 21). *Auditory Brainstem Response*. Retrieved December 21, 2010, from <http://emedicine.medscape.com/article/836277-overview>
- Campbell, K. C. M., & Mullin, G. Emedicine from *WebMD*. (2009, December 18). *Impedance Audiometry*. Retrieved December 20, 2010, from <http://emedicine.medscape.com/article/1831254-overview>
- Campbell, K. C. M., & Mullin, G. Emedicine from *WebMD*. (2009, December 18). *Otoacoustic Emissions*. Retrieved December 21, 2010, from <http://emedicine.medscape.com/article/835943-overview>

8. Kimball, S. H. Emedicine from *WebMD*. (2010, May 11). *Speech Audiometry: Treatment and Medication*. Retrieved December 21, 2010, from <http://emedicine.medscape.com/article/1822315-treatment>

9. Kutz, Jr., J. W., Mullin, G., & Campbell, K. C. M. Emedicine from *WebMD*. (2010, March 31). *Audiology, Pure-Tone Testing*. Retrieved December 21, 2010, from <http://emedicine.medscape.com/article/1822962-overview>

10. Kutz, Jr., J. W., Mullin, G., & Campbell, K. C. M. Emedicine from *WebMD*. (2010, March 31). *Audiology, Pure-Tone Testing*. Retrieved December 21, 2010, from <http://emedicine.medscape.com/article/1822962-treatment>

Google Images retrieved on December 23, 2010 from <http://images.google.com/> and National Center for Hearing Assessment and Management (NCHAM) on December 23, 2010 from <http://www.infanthearing.org>

This fact sheet was reprinted, with permission, from the Nevada Dual Sensory Impairment Project • Spring 2011 Newsletter • Volume 20, Number 1, Pages 3-5

For more information:

Colorado Department of Education

Phone Number: 303-866-6694 – Ask to speak with a Deaf-Blind Specialist on staff with ESSU

Fax: 303-866-6918

Exceptional Student Services Unit

1560 Broadway, Suite 1100

Denver, CO 80202

Web Page Address:

<http://www.cde.state.co.us/cdesped/SD-DB.asp>

Fact Sheets from the Colorado Services to Children and Youth with Combined Vision and Hearing Loss Project are to be used by both families and professionals serving individuals with vision and hearing loss. The information applies to children, birth through 21 years of age. The purpose of the Fact Sheet is to give general information on a specific topic. The contents of this Fact Sheet were developed under a grant from the United States Department of Education (US DOE), #H326C080044. However, these contents do not necessarily represent the policy of the US DOE and you should not assume endorsement by the Federal Government. More specific information for an individual student can be provided through personalized technical assistance available from the project. Reviewed: 2/17